

ADVANCED CARDIAC THERAPIES IMPROVING OUTCOMES NETWORK

HeartMate 3™ LVAD

Patient Education Handbook



action 

action

ADVANCED CARDIAC THERAPIES
IMPROVING OUTCOMES NETWORK

Working together to improve critical outcomes for all heart failure patients.

The Advanced Cardiac Therapies Improving Outcomes Network (ACTION), was established to improve the care of patients with heart failure. ACTION unites all stakeholders (providers, patients/families and researchers), to share experiences, improve education and search for best practices to drive improvement in areas that are often untouched by clinical trials alone.

The learning network approach allows for critical improvements to be made faster across a collaborative system.

Visit our online education site myactioneducation.org (originally designed for pediatric patients and their families) to learn more about the following:

- understanding your heart failure diagnosis
- medicine treatment options
- knowing your device and surgery options
- maintaining your health and wellness

For more information about the learning network approach or the work ACTION is doing, visit actionlearningnetwork.org

This is **MY** VAD Journey

Hi, my name is:



(_ _) _ - _ - _ - _ - _ - _ -



My caregiver is:



Quick References

It's important to keep some basic information about your diagnosis and surgeries available for quick reference.

My VAD Care Team Contact Info



(---) --- - ---



My Hospital

Name

Street Address

City, State, Zip

- My Heart Disease -



I am Allergic to:

♥

♥

♥

♥

♥

My Past Surgeries

My VAD Surgery | Date:

Notes: _____

My driveline is on the:

Left Right

I am listed for transplant:

No Yes _____
Listing Date

My pump parameters: _____

RPM: _____ Power: _____ Flow: _____ PI: _____
Range Range

I can feel my pulse: Yes No



Take my blood pressure using a: _____

My blood pressure goal: _____

Blood thinner medicine(s) I take:

My INR goal:



Using this Handbook

This is a quick educational reference guide and scrapbook to keep track of topics that may come up during your journey. There are times when your care team may ask you to follow different instructions, make sure to write down these important pieces of information.

Additional education will be provided by your care team. If you need more information or clarification on certain topics or questions, please ask your care team.

For complete instructions regarding the HeartMate 3™ LVAD, please refer to your ***HeartMate 3™ Left Ventricular Assist System Patient Handbook***.

©2024 Cincinnati Children's Hospital Medical Center. All Rights Reserved.

The contents of this VAD book, including text, graphics and other materials ("Contents") is offered solely for educational and informational purposes. The information should in no way be considered as offering medical advice for a particular patient or as constituting medical consultation services, either formal or informal. The Contents do not constitute either an explicit or implicit consent or contract by any physician or other employee of Cincinnati Children's Hospital Medical Center or other member of the ACTION Network (collectively, "Network") to create a physician-patient relationship and such information in no way creates, substitutes or in any way consists of examination, diagnosis, treatment or the prescription of treatment for any patient. Content users must use their independent judgment in determining the value and use of the information contained herein and its application to a situation. The Network expressly disclaims any decision-making authority or supervisory control over the Content users. No information provided herein shall create a warranty of any type nor shall any person rely on any such information or advice. Views expressed or implied in the Contents are not the official positions or views of the Network. The Network makes no representations or warranties of any kind, express or implied, about the completeness, accuracy, or reliability of the information contained in the Contents.

IN NO EVENT WILL ANY MEMBER OF THE NETWORK OR THEIR RESPECTIVE TRUSTEES, OFFICERS, EMPLOYEES, AGENTS, VOLUNTEERS, OR REPRESENTATIVES BE LIABLE TO YOU OR ANY OTHER PARTY FOR ANY DIRECT, INDIRECT, CONSEQUENTIAL, EXEMPLARY, INCIDENTAL, SPECIAL OR PUNITIVE DAMAGES OR CLAIMS OF ANY KIND.

Table of Contents



- 1. Introduction to Your HeartMate 3™ LVAD 09**
HeartMate 3 LVAD Patient Journey
What is a VAD?
How will a VAD help me?
Why do I want a VAD?
Who will be my care team?



- 2. Surgery & Hospital Stay 19**
Expectations
Surgery
Intensive Care Unit (ICU)
Transfer Ready
Preparing to Leave the Hospital



- 3. Device Education 27**
Online Resources
HeartMate 3 LVAD System Components
Mathematics & Mechanics
HeartMate 3 LVAD Quiz



- 4. Self Care 41**
Labs
Tests
Medicines
Blood Pressure
Nutrition & Hydration
Weight
Emotional Wellbeing
Driveline Care

Table of Contents



5. Potential Challenges 53

- Infections
- Strokes
- Bleeding
- Hemolysis
- Pump Clots



6. Daily Life and Safety 59

- Your "Go Bag"
- Dental Visits
- Colds & Non-VAD Infections
- Over the Counter Medicines
- Travel
- Return to Daily Life
- Exercise
- Intimacy



7. Your Journey Continues: Resources 67

- The Waitlist
- Reasons to Call Your Care Team
- Keeping You Safe
- Emergency Services & Medical Safety Card



8. Glossary & Appendix 73


- Glossary
- Word Search
- Appendix

Introduction to Your HeartMate 3™ LVAD

- ♥ HeartMate 3 LVAD Patient Journey
- ♥ What is a VAD?
- ♥ How will a VAD help me?
- ♥ Why do I want a VAD?
- ♥ Who will be my care team?


HeartMate 3™ LVAD Patient Journey*

Note: All patients are unique and may not follow the path exactly as shown. Some steps will need to be determined by your care team and/or some may not apply.



What is a VAD?

How does it work?



VAD stands for “Ventricular Assist Device”. It helps a weak or really sick heart to pump blood to the body. The type of VAD that we are talking with you about is a continuous flow VAD, called the **HeartMate 3™ Left Ventricular Assist Device (LVAD)**.

Your HeartMate 3 LVAD will work by continuously spinning blood from the weak part of the heart, through the device, out to the **aorta**, and the rest of the body. There is a small rotor located inside the VAD that helps with this spinning motion. The pump is always connected to a controller and requires a power source to keep the VAD spinning. Because the continuous flow device takes on some or all of the work of the weakened heart chamber, you may not be able to feel your pulse.

Where is the VAD implanted?

The VAD can be **implanted** in any chamber of the heart but usually is placed in the left ventricle. The location of the device determines what your VAD may be called. With your care team, check the box next to the kind of VAD that is right for you. For purposes of this book, your VAD will be referred to as either a VAD or a HeartMate 3™ LVAD



LVAD: A left ventricular device implant, or LVAD, is implanted in the left ventricle of the heart. It improves blood flow to the entire body.



SVAD: A systemic assist device is for patients who were born with only one working ventricle. A VAD is implanted into the “systemic” ventricle that pumps blood to the body.

Implant Location Notes...

How will a VAD help me?

VADs are placed for different reasons. Check the boxes below that apply for why you need a VAD.

- The VAD will help my body prepare for transplant while I wait for the best heart for me to be available.
- The VAD will make me feel better and allow me to spend more time with my family and friends even if a transplant is not right for me.
- The VAD will give more blood flow to my body and make me feel better but my future journey is still unknown.



Why do I want a VAD?

What are your biggest worries about a VAD? What will you hope the VAD will help you do? Use the space below to note any concerns or reasons you think a VAD will be right for you.



A large, rounded rectangular box with a yellow border, containing 20 horizontal lines for writing. The box is positioned in the center of the page, below the introductory text. The lines are evenly spaced and extend across the width of the box. The background of the page features abstract teal and light blue geometric shapes.

Who will be my care team?

There are many people that will care for you during your journey. You and your family are a very important part of the team and will always help to make decisions about your care.

Fill in your care team member's names under their titles.



Cardiac Surgeon _____

Heart Failure Doctor _____

VAD Coordinator _____

Nurse Practitioner _____

ICU Doctor _____

Nurse(s) _____

Physical Therapist(PT)/
Occupational Therapist(OT) _____

Psychologist _____

Cardiac Rehabilitation
Specialist _____

Nutrition Specialist _____

Pharmacist _____

Social Worker _____

Surgery & Hospital Stay

- ▼ Expectations
- ▼ Surgery
- ▼ Intensive Care Unit (ICU)
- ▼ Transfer Ready
- ▼ Preparing to Leave the Hospital

Expectations

Many patients that have VAD surgery notice an improvement in symptoms while they are in the hospital, but it can take months for you to get stronger and recover. As you recover, you may feel sad, have pain and be anxious. This is normal. Be honest about how you're feeling and share your feelings with your care team. Here are some other things to keep in mind:

- ♥ You will have to work hard on your exercises, both in and out of the hospital, in order to get strong (it can take 3–6 months to recover).
- ♥ Medicines and time will help lessen the pain.
- ♥ Sometimes things may not go as planned and there will be bumps in the road. Ask questions if you feel like things are not going as planned.
- ♥ The **Intensive Care Unit (ICU)** can be loud and it may be hard to sleep at times. If you are having trouble sleeping, talk with your team about options to decrease the noise and interruptions.
- ♥ What you eat is really important. Your care team will help you make proper food choices.
- ♥ Your care team will help keep you comfortable and teach you a lot of information in order to keep you safe after surgery.

Changes in Your Journey

There are times when a plan may not go as expected, causing your journey to take a different path. Sometimes patients receiving VADs experience **complications**. If you are on the transplant waitlist these complications may be serious enough to cause your removal from the waitlist. If this happens, your care team will talk to you about the options.

Additionally, if you experience any complications that cause the VAD to not work properly, it may need to be turned off. Your care team will be there with you throughout your journey.

Surgery

Questions I have...	Answers received...
_____	_____
_____	_____
_____	_____
_____	_____
_____	_____

How is the VAD placed in my chest?

The surgery can take an entire day. Your surgeon either performs a **sternotomy** (an incision in the chest bone), or a **thoracotomy** (an incision between the ribs on your left side), and implants the VAD into your heart. This will leave a scar in the middle or side of your chest. The surgery requires a heart and lung bypass machine to circulate blood to your body during the operation. After the surgery your heart and VAD work together to send blood to your body.



My surgery took _____ hours.

How I feel after my surgery...

Intensive Care Unit (ICU)

After the operation, you will recover in the ICU. You will most likely have a breathing tube that's connected to a ventilator (breathing machine) and you will be connected to many lines, tubes, and medical equipment (see illustration below).

This equipment is necessary to monitor your body and give medicine needed for recovery. The equipment and lines can be scary, but are completely normal. Day by day, your care team will work towards removing lines and tubes. The breathing tube is one of the first tubes/lines to be removed. It is also important to get out of bed as soon as possible to help with the recovery process.


1 My breathing tube was removed on _____.

2 My 1st driveline dressing change was on _____.

3 My arterial line was removed on _____.



I got out of bed post-surgery day # _____.



Intensive Care Unit (ICU)


Unplanned Events

Your care team watches closely for bumps in the road or issues that may arise. Some of the issues that can occur during your hospital stay include:

Stroke: With all VADs there is a risk of stroke, which is caused by bleeding or a blood clot in your brain. Both may cause injury to your brain. Your care team manages your medicines carefully to prevent strokes, however sometimes strokes may occur.

Chest Bleeding: After the operation there may be some bleeding from the chest that is captured and removed by the chest tubes. Over time the bleeding will slow and the fluid will become clear. This is when **anticoagulation** medicines (blood thinners) will be started to prevent the pump from getting a clot inside of it.

Infection: When you come out of the operating room you will be on **antibiotics** to prevent infection. Everything will be kept as clean as possible. Your care team will perform frequent **dressing** changes to prevent germs from causing an infection at your surgical site and driveline site.



Right Heart Function: *(for patients with LVADs and two ventricles only)*

For the LVAD to work, the right side of the heart has to be able to move blood to the left side of the heart. When it doesn't, this is called "right heart failure." It's difficult to predict how the right heart will handle surgery. In some instances, you may need to remain on IV medicines or even need a second VAD to help the right side of the heart move blood to the left side of the heart.


Fluid Management: The pump needs *preload*, or a certain amount of blood that enters the heart, in order for the pump to send fluid to the rest of the body. It can be difficult to determine if you have too much water in your body or if you are *dehydrated*. To make sure there is a perfect fluid balance, medicines (*diuretics*) are given to help get rid of extra fluid. There may also be a change in how much you are allowed to drink each day. Sometimes, you will need more fluid for your pump to work well, and your care team will ask you to drink a minimum amount each day.

High or Low Blood Pressure: When your blood pressure is too high, the device will have trouble getting blood to the body. When your blood pressure is too low, the blood may have trouble traveling back to the pump. Your care team will determine a target blood pressure goal and adjust your medicines to meet the goal. Your care team may be measuring your blood pressure differently now that you have your VAD.

Blood Pressure Cuff



Doppler



Transfer Ready

If available at your care center, the Cardiac Floor or Step Down Unit is where you will be transferred after the ICU. These areas of care centers are for patients who are not critically ill but are still recovering after surgery.



After VAD surgery, I was in the ICU for _____ days. I transferred out of the ICU on _____.



Your focus will be on getting stronger. In addition, this is the time that education becomes even more important so that you can be **discharged** out of the hospital.

Preparing to Leave the Hospital

There are specific goals that must be met in order to be **discharged**, or leave the hospital. You and your care team will decide the best places for you to recover from your surgery. This may include going to a **rehabilitation (rehab) center** before you go home. At rehab, the team will focus on helping you rebuild your strength so you can safely go home. There are doctors, nurses, and other therapists there that will continue to monitor you.

Use the **HeartMate 3™ LVAD Patient Journey** on page 11 to follow along and note your specific goals. *NOTE: All patients are unique and may not follow the path exactly as shown. Some steps will need to be determined by your care team and/or some may not apply.*



I was discharged on _____ to

Rehab Center

Home

Other.


Device Education

- ♥ Online Resources
- ♥ HeartMate 3™ LVAD System Components
- ♥ Mathematics & Mechanics
- ♥ HeartMate 3 LVAD Quiz

Online Resources

Learning about your HeartMate 3™ LVAD is an important part of getting you ready to be discharged and keeping you safe. Your care team and our online educational platform: myactioneducation.org will provide more in depth training with electronic modules and videos to help prepare you for the HeartMate 3 LVAD quiz you and your caregivers must complete before leaving the hospital.

The VAD equipment is life-saving and while managing it may feel overwhelming, learning how each of the pieces work together will help you feel more comfortable in your daily journey.




myactioneducation.org

My Login Information

Username: _____

Password: _____




3
Monitor / HeartMate Touch™ Communication System

4
Controller



5
Batteries (with Clips)

2
Driveline




HeartMate 3™ LVAD System Components

Your HeartMate 3 LVAD should always be connected to the system controller and will require two power sources to work. It is important to always have backup equipment with you wherever you go.

1 HeartMate 3 LVAD

The HeartMate 3 LVAD is placed inside your chest. During the surgery, a small hole is cut into your heart muscle and the pump holder (called the apical sewing ring) is attached to the heart muscle.

The HeartMate 3 LVAD is then inserted through the sewing ring and locked into place. The outflow graft is a flexible tube that connects the outflow of the HeartMate 3 LVAD to the aorta. It helps carry the blood from the pump to the rest of the body. The inside of the pump has a rotor that spins continuously and can distribute up to 10 liters of blood per minute (L/min).



2 Driveline

The driveline is made of electrical wires that connect the HeartMate 3 LVAD to the system controller. The driveline has two important functions in order to make the device work properly. First, the driveline brings power to the HeartMate 3 LVAD to make the pump spin. If the driveline is ever disconnected from the system controller your HeartMate 3 LVAD will stop working. Secondly, the driveline sends important information back and forth between the HeartMate 3 LVAD (inside the heart) and the system controller.




There are two parts of the driveline, the pump cable and the modular cable. The pump cable is the section that is mostly inside the body. The modular cable is the section that connects the pump cable to the system controller through a metal connection. If the modular cable is damaged, it can be replaced without having to exchange the pump.

HeartMate 3™ LVAD System Components

3 System Monitor / HeartMate Touch™ Communication System

You will only be connected to the monitor when you are in the hospital or in clinic. The monitor tells your care team how your device is working so they can make any adjustments needed.



4 System Controller

- (A) **Cable Disconnect Symbols:** light up if a power cable or driveline has been disconnected
- (B) **Battery Button:** shows how much battery power remains
- (C) **Pump Running Symbol:** stays on (and green) to show that the pump is working
- (D) **Display Button:** allows the user to advance through settings and alarms
- (E) **User Interface Screen:** displays the VAD settings and other important information
- (F) **Alarm Symbols:** the red broken heart and wrench symbols notify when there is a problem with the controller or pump, and the battery symbol notifies when the batteries are disconnected or have less than 5 minutes of power remaining

- (G) Silence Alarm Button:** silences active red alarms and power cable disconnected alarm for 2 minutes, low battery yellow diamond alarms for 5 minutes and yellow wrench alarms for 4 hours
- (H) Battery Status Symbol:** displays how much battery power remains when the battery status button is pressed
- (I) Driveline Connector:** the cable connecting the driveline to the controller
- (J) Power Cable Connectors:** attach to the power module, mobile power monitor module or batteries and provide power to the controller

The driveline connects to the controller through the **driveline connector (I)**. The system controller has **two power cables (J)** that connect to a power source. There is a backup battery in the system controller that powers the controller for approximately 15 minutes in the event that the power becomes disconnected. This is for emergency use only. You will have a primary and a backup system controller. Always carry the backup system controller and spare batteries in the event that you need to change them quickly.

The system controller has a display screen **(E)** that provides VAD settings: Speed (RPM), Flow (L/min), Power (watts) and Pulsatility Index *(See Math & Mechanics section for more information)*. You can see each setting by pressing the display button. If there is an active alarm, the screen will provide information and the system controller will have lights to alert you.

Reminder: complete a self-test daily by holding the battery button for 5 seconds. All the lights turn on and alarms will sound for 15 seconds and then they will turn off. If this does not occur, follow directions from your care team.




HeartMate 3™ LVAD System Components

5 Batteries (with Clips)


A pair of batteries can last **10–17 hours**. The battery indicator on the system controller (**H**) will show how much power a battery has remaining. The battery indicator has four bars. Each bar represents about 25% of the battery life remaining. If the battery is less than 15 minutes, the diamond indicator will turn yellow. The battery indicator will turn red if there is less than 5 minutes remaining.

After checking for full charge, connect the batteries to the clip by lining up the red arrows and inserting the battery into the clip until you hear a click.

To connect a power cable to the clip, align the prongs to make a complete connection. Firmly push the power cable directly to the battery clip. Twist the connector clockwise to tighten the connection. When disconnecting the battery, untwist the connector and pull directly out from the clip.



Batteries with Clips



Battery Charger

Battery Charger

The battery charger is a charging station for up to four batteries. It takes **4 hours** to completely charge each battery. The status symbol will turn green when a battery is fully charged. Check the symbol before connecting a battery to the controller. If the status symbol is flashing red, do not use the battery and it must be replaced.

Mobile Power Unit

The Mobile Power Unit (MPU) plugs into the wall and should be used whenever you are sleeping. Connect to your batteries when you are up moving around to avoid build-up of static electricity. The MPU has a black cord that plugs into a standard wall power outlet. Make sure that this connection is always tight. The power cables connect to the patient cable the same way that the battery clips are connected. Always connect black-to-black and white-to-white. In the event of a power outage, quickly connect to battery power or a different power source.

Alarms

Alarms can be loud and scary but they are meant to let us know when your HeartMate 3™ LVAD and/or system controller is not working properly. They also can let your care team know if there is a change in how your heart is working. There is no need to panic when you hear an alarm because there are several steps that you can take to correct the problem. When you hear an alarm, check the display screen to determine what is wrong. Use the troubleshooting guide below or refer to the HeartMate 3 LVAD alarm guide for more information on alarms. **Notify your care team immediately if you experience any of the critical alarms listed below.**

ALARM TYPE	SYSTEM CONTROLLER MESSAGE/SYMBOLS	REASON	ACTION TO TAKE
CRITICAL ALARMS <i>Contant Tone Sounded</i>			
Pump Off	"Call Hospital Contact" AND "Low Flow" alternate AND  <i>flashing red</i>  <i>solid black</i>	Pump has stopped running, possibly because power has been disconnected or failed.	<i>Immediately connect to a power source (if disconnected/failed).</i> <i>If restoring power does not resolve, press any button on the Controller to attempt pump start, and immediately call care team.</i>

[more >](#)

HeartMate 3™ LVAD System Components

ALARM TYPE	SYSTEM CONTROLLER MESSAGE/SYMBOLS	REASON	ACTION TO TAKE
No External Power	<p>“Connect Power Immediately” and Backup Battery graphic alternate AND</p> <p> <i>flashing red</i></p> <p>) <i>flashing yellow</i> (next to power cables)</p>	<p>The Controller is not receiving power from either power cable.</p> <p>The pump is being powered by the Controller’s backup battery.</p>	<p><i>Immediately connect to a working power source (Mobile Power Unit or two charged batteries) to ensure pump does not stop.</i></p>
Pump Stop AND No External Power	<p>“Connect Power Immediately” AND “Backup Battery” alternate AND</p> <p> <i>flashing red</i></p> <p>⌚ <i>solid black</i></p> <p>) <i>flashing yellow</i> (next to power cables)</p> <p> <i>flashing red</i></p>	<p>Pump has stopped running and mobile power unit has failed, possibly due to static electricity.</p>	<p><i>Immediately connect to charged batteries.</i></p> <p><i>If restoring power does not resolve, press any button on the Controller to attempt pump start</i></p>
Driveline Disconnected	<p>“Connect Driveline” flashes</p> <p> <i>flashing red</i></p> <p>) <i>flashing red next to driveline connector</i></p> <p>⌚ <i>solid black</i></p>	<p>Driveline is disconnected from the Controller or the connector is broken.</p>	<p><i>Reconnect the Driveline to the Controller. Move Driveline Safety Lock on the Controller to the locked position. Make sure Modular inline Connector is secure.</i></p> <ul style="list-style-type: none"> - <i>If alarm persists after reconnection, press any button on the Controller to attempt pump start.</i> - <i>If Driveline alarm still persists, replace Controller with backup.</i> - <i>If alarm still persists, call care team.</i>
Low Flow	<p>“Call Hospital Contact” AND “Low Flow” alternate AND</p> <p> <i>flashing red</i></p>	<p>Pump flow is less than 2.5 lpm</p>	<p><i>Make sure the Driveline is connected to the Controller and a power source is connected to the Controller. Call care team for diagnosis.</i></p>
Low Battery (less than 5 min)	<p>“Low Battery” AND “Replace Power Immediately” alternate AND</p> <p> <i>flashing red</i></p>	<p>Less than 5 minutes of battery power remains.</p>	<p><i>Immediately connect to a working power source (Mobile Power Unit or two charged batteries).</i></p>

ALARM TYPE	SYSTEM CONTROLLER MESSAGE/SYMBOLS	REASON	ACTION TO TAKE
Controller Hardware Fault	“Call Hospital Contact; Controller Fault” AND All Symbols are off AND All Controller buttons are non-functional	Controller is not functioning. Pump will operate as long as power is applied to the controller and no other malfunction occurs. All alarms are not functional.	<i>Call care team immediately for diagnosis and instructions.</i> <i>Switch to backup Controller if instructed.</i>

System Controller Change Out

The HeartMate 3™ LVAD cannot run without the system controller. If the system controller is not working, the HeartMate 3 LVAD will stop spinning and may make you feel dizzy, or may even cause you to faint. Stay calm, sit or lay down and contact your care team. If instructed to do so by your care team, have someone quickly get your backup controller to perform a system controller change out. Once the backup controller is located, place it in front of the faulty controller and follow these steps:

- 1 Connect the backup batteries to the clips (skip this step if using the mobile power unit).
- 2 Connect the backup controller to a power source (batteries or mobile power unit).
- 3 Open the safety lock on the damaged controller and press the red button to release the driveline. While pressing the red button, pull the driveline out of the damaged controller.
- 4 Line the driveline arrow to the controller arrow and insert the driveline into the new controller until you hear a “click.” Check that the double arrow symbol is green indicating the pump is turned on.

HeartMate 3™ LVAD System Components

My HeartMate 3™ LVAD Checklist



Scan QR code to
download checklist.

Daily

- ♥ Perform a system controller test—hold battery button until alarm rings and symbols light up
- ♥ When removing batteries from charger check battery power before connecting to system controller
- ♥ Check that the arrows on the system controller are green
- ♥ Scroll through VAD numbers and write them down
- ♥ Check for the green light on the mobile power unit before plugging in at nighttime
- ♥ Check to make sure the safety lock is covering the red release button

Weekly

- ♥ Complete driveline dressing care as instructed
- ♥ Check the modular connection and driveline for any damage
- ♥ Check the battery charger is free of dust and debris
- ♥ Rotate the unused batteries in the charger slots

Monthly


- ♥ Inspect and clean the battery and battery clip contacts with a cotton swab dampened with alcohol
- ♥ Check for any damage to connections
- ♥ Check the shower bag for any damage

6 Months – Yearly

- ♥ Replace the AA batteries in the mobile power unit
- ♥ Inspect both system controllers for damage
- ♥ Your care team will help you recalibrate all batteries. Check battery expiration date—if it has been used for more than 3 years, contact your care team
- ♥ Have your equipment serviced by your care team. Connect the back up controller to a power source and allow it to charge completely. Afterward, perform a self test on that controller.

Mathematics & Mechanics

What Can Your Care Team Change on the Device?



Pump Speed or Revolutions Per Minute (RPM):

How fast the rotor spins is measured in RPM. This is important because the speed determines how much blood can flow through the device to the body. The speed is the only setting your care team can change.

What Does the Device Calculate and Display?



Cardiac Output (L/Min): The amount of blood that flows through the device and is pumped to the body is measured in “liters per minute (L/min).”



Power (Watts): Amount of “work” the pump is doing to get blood to move through the body. How much power the pump is using is measured in “watts.” The controller and power source provides power to the pump to keep it spinning.



Pulsatility Index: Provide details on how the pump is functioning – these numbers help guide fluid and blood pressure management.

What Does Your Care Team Calculate?



Cardiac Index (L/Min/m²): Amount of blood flow moving through the body in relation to the size of the patient, also known as body surface area (BSA)*.

**Body Surface Area (BSA)= takes into account weight & height*

HeartMate 3™ LVAD Quiz

Once your education is complete, you will need to complete a quiz about your equipment. The quiz is to make sure you are comfortable with the equipment.

Ask your care team for the web link to complete the HeartMate 3 LVAD quiz.



myactioneducation.org

My quiz is available at:

Self Care

- ♥ Labs
- ♥ Tests
- ♥ Medicines
- ♥ Blood Pressure
- ♥ Nutrition & Hydration
- ♥ Weight
- ♥ Emotional Wellbeing
- ♥ Driveline Care

Labs

You will have laboratory testing (labs), including blood draws, both at scheduled times and whenever your team feels they need to follow you closer. Requested labs may include the following:


ANTICOAGULATION	
Heparin Level or Anti-Xa	♥ measures how thick the blood is when on Heparin
PT/INR	♥ measures how thick the blood is when on Warfarin
PTT	♥ measures how thick the blood is when on Heparin or Bivalirudin
LDH	♥ shows if red blood cells are breaking, signaling there may be a clot in the device
CARDIAC	
BNP/ NT-pro BNP	♥ helps monitor fluid status and how well the heart and VAD are working together
HEMATOLOGY	
CBC	♥ measures red blood cells, white blood cells and platelets in the blood
KIDNEY	
Renal	♥ monitors <i>kidney</i> function and measures fluid status and electrolytes
LIVER FUNCTION	
Hepatic Profile	♥ measures how well the <i>liver</i> is functioning
INFECTION & INFLAMMATION	
CRP	♥ measures if there is any infection or inflammation within the body

Tests

You will have tests at scheduled times and additional ones can be requested whenever your care team thinks you need closer attention. You may have had one or more of these tests before your VAD was placed. Tests include:




CT Scan: If the VAD is acting unusually or the LDH is high the team may order a CT scan of the heart. A CT scan of the head may be performed if there is a concern for stroke.



Echocardiogram: An “Echo” is an ultrasound that uses nonradioactive, high-frequency sound waves to view the heart. An Echo is a non-invasive test to help the VAD team diagnose any heart or VAD problems.



Exercise Test: To determine how well you are doing on the VAD, your care team may order an exercise test. An exercise test may be performed on a stationary bike, a treadmill or as a six minute walk.



Ramp Study: A ramp study is performed to determine the best speed (RPM) for your VAD device to be set to. Your care team will watch how your heart reacts as the speed of the device is changed. The ramp study can also be performed if there is concern for a blood clot in the device.



IMPORTANT Once you have a VAD, **DO NOT have an MRI test.** A MRI uses strong magnets that attract metal objects and would damage your VAD.

Medicines

In the hospital, medicine will be given through an IV. Your care team will change those medicines to be given by mouth when preparing for discharge. It's important the medicines are taken at the same time every day. By the time you are ready to go home, medicines will likely be taken twice a day. Use a pill box to keep your medicines organized. Inform your care team if you take any additional medicines NOT prescribed by them.

Below is a list of medicines you may be taking when discharged. Check off the medicines you are taking and write in any additional ones that are not listed below.

- | | |
|---|---|
| <input type="checkbox"/> Amlodipine | <input type="checkbox"/> Sildenafil (Revatio) |
| <input type="checkbox"/> Antibiotics | <input type="checkbox"/> Spironolactone (Aldactone) |
| <input type="checkbox"/> Aspirin | <input type="checkbox"/> Warfarin (Coumadin) |
| <input type="checkbox"/> Chlorothiazide (Diuril) | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Clonidine | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Dapagliflozin (Farxiga) | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Enalapril/Lisinopril | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Entresto | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Furosemide (Lasix) | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Hydralazine | <input type="checkbox"/> _____ |
| <input type="checkbox"/> Omeprazole/Pantoprazole (Protonix) | <input type="checkbox"/> _____ |

Medicines

Blood Thinners

When you are on a VAD, you will usually be on blood thinner medicine. Your care team will increase and decrease the medicine based on your lab results.

Aspirin: You may take aspirin to prevent platelets from sticking together and forming a clot in your VAD.

Warfarin (Coumadin): Once your care team determines the time is right, you may take warfarin pills by mouth. Warfarin is the most important medicine you will take, but it can be difficult to get your dose (the amount you take) correct. The dose needed will go up and down frequently depending on your INR and it can take a number of days to get it right. Once your dose is correct, your IV blood thinner medicine will be turned off.

Warfarin is what makes your INR (blood levels for anticoagulation or how thick or thin your blood is) increase. Below are causes for:



High INR (*thin blood*)

- ♥ New medicines
- ♥ Prolonged vomiting/diarrhea*
- ♥ Prolonged inability to eat*
- ♥ An increase in alcohol consumption
- ♥ A **decreased** intake of food or drink containing vitamin K

**Discuss with your care team*



Low INR (*thick blood*)

- ♥ Missed doses of warfarin
- ♥ An **increased** intake of food or drink containing vitamin K
- ♥ Large increase in exercise




My goal INR is: _____

All warfarin tablets (no matter the brand name) have a color and shape that indicate their strength, or how many milligrams (mg) are in each tablet. There may be frequent changes in your warfarin dosing, please be familiar with your dose's tablet color and shape. Contact your care team ahead of time if you need refills on medicine so you don't miss taking any of your doses.

1 mg	2 mg	2.5 mg	3 mg	4 mg	5 mg	6 mg	7.5 mg	10 mg
Pink	Lavender light purple	Green	Tan	Blue	Peach light orange	Teal blue-green	Yellow	White

A **consistent diet** is very important while taking warfarin. Warfarin blocks the ability for the body to activate vitamin K, which helps the body produce blood clots. When vitamin K interacts with warfarin, it can cause changes in your INR level. If you eat foods rich in vitamin K, such as leafy green vegetables, you will need more warfarin to keep your blood thin. *Note: Other medicines can also affect how well the warfarin works.* Below are some examples of vitamin K rich foods from high to low to be mindful of while on the medicine. Foods with high to medium amounts of vitamin K will interact with Warfarin the most.



information above completed on ____/____/____

Medicines

High Blood Pressure

At times, on a VAD you may have high blood pressure, which will prevent the VAD from pumping well and can increase your risk of stroke. Blood pressure medicine examples include:

- ♥ Enalapril/Lisinopril or Entresto
- ♥ Amlodipine
- ♥ Clonidine
- ♥ Other blood pressure medicines

Preventing Infection

Antibiotics are given in the operating room and ICU to prevent infection. They are also given if your driveline gets infected. You may need antibiotics indefinitely to prevent infection. Antibiotics can make your INR high or low.

Immediately notify your care team if you start taking an antibiotic.

Stomach Ulcers & Acid Reflux

Medicines to prevent “heartburn” are used to decrease the acid in the stomach. This may help with decreasing your stomach discomfort.

Examples include:

- ♥ Omeprazole/Pantoprazole (Protonix)
- ♥ Lansoprazol (Prevacid)

Extra Body Water (Diuretics)

Diuretics were important when you were in heart failure. When you are on a VAD you should require less, but you still may need a small amount to get the fluids in your body just right. Examples include:

- ♥ Furosemide (Lasix)
- ♥ Chlorothiazide (Diuril)
- ♥ Spironolactone (Aldactone)

Blood Pressure

Controlling your blood pressure is important. Your care team will determine a target blood pressure goal and adjust medicines to meet it. Blood pressure can be monitored using a blood pressure cuff and/or a Doppler. This depends on if you can feel your pulse. If you go to an outside hospital or ride in an ambulance you may need to tell them the best way to take your blood pressure.



My blood pressure goal is: _____.

The best way to take my blood pressure is using a: (Check one)



Blood
Pressure
Cuff



Doppler



Nutrition & Hydration

Once you have your VAD you must stay on a stable diet. Eating a healthy diet will help you heal and get stronger.



Fluids

To function properly, the VAD needs a certain amount of fluid flowing through it. Preload is the amount of fluid in your body, or how “full” the heart is. Dehydration from not drinking enough, or increased fluid losses

Nutrition & Hydration

(vomiting, diarrhea, sweating) will lead to low preload. High preload can be caused by heart failure or not peeing enough.



If you have **low pre-load** (dehydration) you may experience:

- ♥ lightheadedness, tiredness, falls
- ♥ alarms from your VAD

If you have **high preload** (fluid-overload) you may experience:

- ♥ swollen face, hands or legs
- ♥ poor appetite
- ♥ shortness of breath

VAD speed and flow may need to be changed over time as your fluid balance changes. It's important to track your weight and fluid intake, so your care team knows if your fluid balance is off.



My goal is to drink _____ of water per day.

Weight



My weight at discharge is _____ lbs/kgs.




I will weigh myself at home every _____ day(s).



Contact my care team if my weight is:

above _____ lbs/kgs

below _____ lbs/kgs



Emotional Wellbeing

For some patients, having a VAD and needing medical care can be hard. You or your caregiver may feel sad, worried, or even angry at times. Taking care of your emotional and mental health is a very important part of your care. Be sure to talk with your care team if you have these feelings or any changes in behavior. Your care team will be able to connect you with the right support services. Your center may have a patient and family VAD group you can join. Abbott also has a patient ambassador program which can help connect you to others in the HeartMate™ LVAD Community. You can join online at: abbott.com/ambassadors.

Things that make me feel better when worried or upset:

Driveline Care

The driveline connects the VAD on the inside of the body to the system controller on the outside of the body. If an infection develops on or around your driveline site, it can be very serious. Depending on how bad it is, the infection may need to be treated with IV or long-term antibiotics taken by mouth, or it may require being admitted into the hospital.



It's important to properly care for the driveline site to prevent infection. Germs that collect at the driveline site could travel to the heart if left untreated.

Frequent movement of the driveline can cause damage above

and underneath the skin and increase the risk of infection. Using anchors to keep the driveline in place is important to keep the driveline site healthy.

Dressing Changes: Your driveline site will need to have dressing changes using *sterile* technique. Everyone should use sterile gloves and wear masks during dressing changes. How often you change your dressings will be determined by your care team. Some patients learn to do their own driveline dressing changes and others have a dedicated caregiver do them. Your care team will give you or your caregiver step-by-step instruction on how to correctly perform sterile dressing changes with your specific dressing kit.



My dressing should be changed every _____ day(s).

Potential Challenges

- ♥ Infections
- ♥ Strokes
- ♥ Bleeding
- ♥ Hemolysis
- ♥ Pump Clots



Infections

If an infection develops on or around your driveline, it can be very serious. Most infections can be treated with antibiotics and changes to your dressing care.

If the infection becomes more severe and spreads to your blood it is called **sepsis** or bacteremia. This will need to be treated with antibiotics through an IV and you will potentially need to be hospitalized. While it is rare, if the infection can't be cleared from the blood with the VAD in place, the device may need to be taken out and a new one put in place, which would require another surgery.

If you notice any changes in your driveline or if your driveline gets pulled, tell your care team right away so it can be treated quickly. Signs and symptoms of a driveline infection include:

- ♥ Redness
- ♥ Pain or tenderness
- ♥ Drainage (new or increased)
- ♥ Swelling and warmth
- ♥ Fever




Strokes

With all VADs there is a risk of stroke, which is caused by bleeding or a blood clot in your brain. Both may cause injury to your brain. Your care team manages your medicines carefully to prevent strokes, however sometimes strokes may occur.

Notify your care team or call 911 if you have any of the following symptoms:

- ♥ Headaches that are different than usual
- ♥ Numbness and/or tingling on one side of the body
- ♥ Weakness on one side of the body
- ♥ Losing feeling and/or movement in the legs or arms
- ♥ Slurring words or trouble when talking
- ♥ Facial expressions and movements don't match (or mirror) on each side of the face
- ♥ Pupil (the dark circle in the middle of the eye) sizes don't match



Bleeding

When on blood thinners, you are always at risk of bleeding. You may experience bleeding from your gums when brushing your teeth and your cuts may bleed more than normal.

Women may experience heavier bleeding during menstrual periods and may need to seek additional treatment.

If you are involved in a traumatic accident, such as a car accident, it can be harder to stop the bleeding. Your care team will need to help.

Nose Bleeds: Blood thinners often can lead to nose bleeds. To prevent nose bleeds, use petroleum jelly or saline nose spray in your nasal passages during cold, dry weather. Do not pick your nose as this may start a nosebleed. If you have a nose bleed:

- ♥ Stay calm. Look forward. Do not tilt your head back.
- ♥ Hold pressure at the bridge of your nose for 10 minutes without letting go.
- ♥ If pressure doesn't stop the bleeding, your care team may advise you to use Afrin® or saline nasal spray.
- ♥ Let your care team know if you can't stop the bleeding, or if you have frequent nose bleeds.
- ♥ Your care team may need to decrease your blood thinner medicine, or you may need to see a special Ear, Nose, and Throat (ENT) doctor.



Gastrointestinal (GI) Bleeds: You may have a more serious bleed in your stomach. Signs and symptoms of a GI bleed include:

- ♥ Stomach pain and no appetite
- ♥ Vomit with blood in it
- ♥ Black, tarry or sticky stools
- ♥ fatigue and shortness of breath

Hemolysis

Red blood cells are a type of cell that circulate in your blood, bringing oxygen to your tissues and organs. When your red blood cells break down, it's called hemolysis. Hemolysis may occur for multiple reasons, the most concerning being a mechanical problem with the VAD. If a clot develops in the VAD it can affect the flow, causing red blood cells to break down as they pass through it. In return, this causes an elevation in your LDH (a marker in your blood of broken red cells) blood test.

Notify your care team if you have any of the following symptoms:

- ♥ Pink, red, cola, or tea colored urine
- ♥ Increased pump power/flow that is out of your "normal" range
- ♥ Decreased pump power/flow that is out of your "normal" range



Pump Clots

Your VAD could develop a clot inside of it, causing it not to work properly. If this happens, and changes to your blood thinner medicine doesn't fix the problem, you may need to get a new VAD. The following symptoms should be discussed with your care team:

- ♥ Pink, red, cola, or tea colored urine
- ♥ Heart failure symptoms such as shortness of breath, exhaustion or abdominal pain/vomiting
- ♥ Increased pump power/flow that is out of your "normal" range
- ♥ Decreased pump power/flow that is out of your "normal" range and low flow hazard alarms



Daily Life & Safety

- ♥ Your “Go Bag”
- ♥ Dental Visits
- ♥ Colds & Non-VAD Infections
- ♥ Over the Counter Medicines
- ♥ Travel
- ♥ Return to Daily Life
- ♥ Exercise
- ♥ Intimacy

Your "Go Bag"

Always have backup equipment with you in case something happens to one of your VAD components. It should not be left in the car during hot summers and cold winters. The following items should always be with you:



- ♥ Emergency VAD contact information
- ♥ Extra controller
- ♥ Extra batteries and clips
- ♥ Water bottle
- ♥ Medicines (needed while away)

♥ _____	♥ _____
♥ _____	♥ _____
♥ _____	♥ _____

Natural Disasters

It's important to have an emergency preparation plan during times of crisis.

Where will I go: _____
NAME

ADDRESS

PHONE

What I will need to take:

- | | |
|-------------------------------|---------|
| ♥ a month supply of medicines | ♥ _____ |
| ♥ dressing supplies | ♥ _____ |
| ♥ equipment | ♥ _____ |
| ♥ go bag | ♥ _____ |

Dental Visits

Good dental *hygiene* is very important. If you get an infection in your mouth it could lead to an infection in your VAD. Before you go to the dentist for a cleaning or other dental work you will need to take antibiotics, called an antibiotic prophylaxis.



My care team wants me to take: _____

Colds & Non-VAD Infections

When you have a VAD you can still get common colds or other infections. Do everything you can to avoid getting sick. Wash your hands frequently, avoid contact with others who are sick, and get a flu shot every year.



Always let your care team know if you get sick. They may direct you to your primary care doctor or ask to see you themselves. What you think may be a common illness could be something more serious for you. Being sick will affect your INR and your care team may need to make changes to your blood thinner medicine. Remember to maintain hydration and nutrition especially when you are sick. If you are unable to do this, contact your care team.

Over the Counter Medicines

Always check with your care team before starting any new prescription, over the counter medicine, or any other medicines or remedies.

When on a VAD, you may have pain, especially after your surgery.



You CAN'T take NSAIDS

(Motrin, Advil, Ibuprofen)
because it changes how
your blood clots



**You CAN take
Acetaminophen**

(Tylenol) for pain

Safe over the counter medicines for me to take include:



Travel

As a precaution, always carry your VAD emergency information with you.

Driving Precautions

Every state has different rules about driving with a VAD, discuss this with your care team. If you are a licensed driver, use caution within the first 90 days after surgery because you may still be taking medication that may affect your concentration. Some medicines will cause side effects that you won't have adjusted to yet.

When you start driving again, take short supervised trips to get used to driving with the new equipment. If you feel dizzy, short of breath, or have blurred vision, pull over to the side of the road immediately and call for help.

Long Distance and Air Travel

You will be able to travel, even on a plane but it takes some extra planning with your care team to make sure you're safe. You will need a travel plan, and an emergency action plan for long-distance trips.

Your care team requests _____ days notice before your trip, so they can make safe plans for you and contact other hospitals to discuss your care.



When traveling by plane, plan for extra travel time when going through the airport. You may also need documentation as you will **not** go through standard security. You will need to tell the airline agents you have a medical device and they will take you through a separate security lane. Make sure to pack all medicines and backup equipment in your carry-on luggage.



Return to Daily Life

It will vary from hospital to hospital, but once your care team decides it's safe for you to return to daily life, multiple steps will occur to ensure your safety.

When you do leave the hospital, make sure to:

- ♥ Train people close to you so they are available to help with alarms and emergencies
- ♥ Notify the local EMS department (fire station, police, etc.)
- ♥ Have backup batteries, controller and any additional information your team gives you to prepare for leaving the hospital

If your care team feels you must stay in the hospital for a longer period of time, discuss how you can continue to progress in your daily care activities.



Sternal Precautions

If your chest incision was made down your sternum (center of your chest) your care team will have specific **sternal precautions**. Ask your team when you can safely:

- ♥ Lay on your stomach/chest
- ♥ Push, pull or lift more than 10 pounds
- ♥ Drive and ride in the front seat

You and your care team will talk about your goals and interests once you have healed from your surgery. This may include returning to work or resuming certain hobbies. It is important to share your goals with your care team.

What I would like my care team to help me do:

Pets

Do you live with a pet? Ask your care team about any precautions you may need to take to stay safe.

Questions about how to safely care for my pet:

Exercise

Once you've recovered from surgery you should be exercising to get stronger. **You will not be able to swim or play contact sports.** You may be instructed to do special exercises at home or you may be asked to come to the hospital frequently to exercise in the clinic. Your care team will help you choose exercises that you can do safely. With your care team, check off the activities that you can do:

Ride a bike (wear a helmet)

Walk or Jog

Tennis/Pickle Ball

Throw a football

Bowling

Yoga

Golf



If you have a wearable tracking device, talk to your care team about coming up with a step goal to keep you moving daily.



My daily step goal is _____ steps.

Intimacy

Your care team will let you know when it is safe to be physically intimate. Talk to your care team about how long you should wait. Your chest bone will need to heal. It is also important to ensure your driveline and system controller are in a secure place to prevent any injury to your exit site. Your care team will have other information to help you feel comfortable returning to this part of your life. They may also talk with you about birth control and family planning choices. Some people find it embarrassing to talk about this part of their life, but it is important to have these discussions so you can return to your normal life safely.

Menstruation

Women who experience monthly periods may see a change in the pattern and amount due to their VAD and the blood thinners. It is important to talk about these changes with your care team. Your care team will review with you when to call or seek emergency care.

Personal questions I have for my care team:

Your Journey Continues: Resources

- ♥ The Waitlist
- ♥ Reasons to Call Your Care Team
- ♥ Keeping You Safe
- ♥ Emergency Services & Medical Safety Card

The Waitlist

With heart failure, patients often feel uncomfortable, sad, frustrated and even angry. Our goal is to help you feel better and get back to doing the things you enjoy. A VAD should help to relieve the symptoms of heart failure, although it may take a while to see the benefit. Always discuss how you feel with your care team so they can help you achieve the best quality of life possible.

If you are on the waitlist for transplant and want more information about how organs are matched, see more from the Organ Procurement and Transplantation Network at: tinyurl.com/3hs6bv9x

Reasons to Call Your Care Team

If you are discharged to go home, you will have a detailed communication plan. The plan will include a phone number to call. Your care team will want to hear about most everything. Examples include:

- ♥ Change in pump parameters, equipment issues and alarms
- ♥ Changes to driveline site
- ♥ Blood pressure outside your goal ranges
- ♥ Changes in your diet
- ♥ New medicines
- ♥ Pain or Bleeding
- ♥ Fever
- ♥ Red, pink, cola or tea colored urine
- ♥ Changes in your mood or anything else you feel uneasy about



You will also have many follow-up clinic visits that are important for you to attend.

Keeping You Safe

When on a VAD you will feel better and you will want to be active. Your care team will talk with you about what is safe and what is not safe.

You CAN'T

take a bath, swim, vacuum or play contact sports.



You CAN

shower (with a shower kit), dance, jog, travel and drive.





Prevent exposure to static electricity when possible. If you are concerned about exposure, make sure to use battery power. Use dryer sheets when doing laundry, wear rubber sole shoes/slippers indoors, and be careful with fleece.



Protect your controller and batteries when it is raining outside or around large amounts of water.



Always wear a seatbelt in the car.



Never disconnect your driveline. Your pump will stop. Never disconnect from both power sources.



Wash your hands frequently and always before dressing changes. This will help keep you healthy.



Always bring your backup equipment when leaving the house. Check the battery charge level before connecting to batteries.



If there is a power outage, switch to battery power. Notify your care team if the power is out for more than 24 hours.



For bleeding cuts, firmly press on the cut for 5–10 minutes, or until the bleeding stops. Call your care team if the bleeding does not stop.

Emergency Services & Medical Safety Card



Local Emergency Services (Medical/Fire/Utilities): Either you or your care team members should notify your local emergency medical department when you are discharged.

It is nice for them to know you are at home with life-saving equipment. If you lose power for more than 24 hours you must go to a new place with power or a generator, or contact your care team if you have questions about what to do.

Medical Safety Card: Always carry your medical safety card with you in case of an emergency. If you don't have a medical safety card, ask your care team to provide the ACTION medical safety card (sample below) for you. You can also download and fill out the card by scanning the QR code to the right.



Scan QR code to download medical Card

VAD Emergency Contact:

PHONE _____

HOSPITAL NAME _____

STREET ADDRESS _____

CITY, STATE, ZIP _____

Sample

action
actionlearningnetwork.org

First Name MRN: _____

DIAGNOSIS: _____

NOTES: _____

Sample

Glossary & Appendix

- ▼ Glossary
- ▼ Word Search
- ▼ Appendix

Glossary

Antibiotics: Medicines to treat bacteria and germs.

Anticoagulation: Medicines to make your blood thinner so your VAD pump does not clot.

Aorta: The large artery that brings blood from the heart to the rest of the body.

Clot: When red blood cells stick together to form a 'glob' in one of your blood vessels. Similar to a scab.

Complications: A disease or problem that happens in addition to the first disease.

Consistent Diet: Always eating the same kind of foods.

Dehydrated: When you do not drink enough water, or you have diarrhea, the volume of fluid in your body goes down. This can make your pump not work as well.

Diagnosis: Determining the cause of the disease or problem.

Discharged: When you leave the hospital and go home, or go to a local place or a hotel.

Diuretics: Medicine to make you pee more.

Dressing: A bandage around the driveline that is coming out of your stomach.

Hygiene: Keeping yourself clean to prevent infection.

Implant/Implanted: During surgery your VAD is placed in your heart.

Intensive Care Unit (ICU): A place in the hospital where patients recover after surgery when they leave the operating room.

Kidney: The organs in your body that remove waste and make urine.

Liver: The organ in your body that cleans the blood.

Preload: The amount of fluid that travels through your blood vessels to your heart.

Rehabilitation (Rehab) Center: A place to go to get stronger and feel better.

Sepsis: An infection in your blood which can make you very sick.

Sterile: Keeping a wound clean. Using a procedure that includes clean gloves and masks to make sure germs do not get near the driveline.

Sternal Precautions: Instructions to follow in order to allow the incision in your chest bone to heal.

Sternotomy: An incision in the chest bone that is made by the surgeon to get to your heart.

Stroke: When a blood clot travels to your brain through a blood vessel.

Thoracotomy: An incision between the ribs on the left side of your chest.

Word Search

W Z X A N A C V
 N L W N N L V M L O V D
 G L O T W S G U H N M F M T A T
 Q W B J I D R M N E B C Y Y T P V O O V
 D E T E I D T N E T S I S N O C L R L L Q E
 C E R Y F F V G A D V D V H J I P I C E C M J Z
 C L S E A H Q O T L I I N S E N L C E R G G T I
 R I T V S B U Z G A U I L C K B S A U P T V Z F
 N R R A P S Z B G N A G M I T U R T L A Q H E E
 X E O X T N I N M D K A A P E N E I G Y H W P D
 E T K Y G R O N T V Q A T O L J O O N A C I E L
 S E R G S O V G C C J Q R C A C N K W K T P
 Z E Q I Y I A W S N T O U R I N S A A A Z B
 X S P D I S C H A R G E D R T T X R J F
 W A N T I B I O T I C S O Z N D C E
 S S U C I L Q Q K X E Y E Y A Z
 I W X G P W B I L I W H W V
 S H F G U G D U X E J T
 P F J M D N I D Q Y
 E Q A P E C V S
 S K B Y Q K
 S T N B
 F U

- | | | |
|--|-------------------------------------|-------------------------------------|
| <input type="checkbox"/> Antibiotics | <input type="checkbox"/> Diagnosis | <input type="checkbox"/> Kidney |
| <input type="checkbox"/> Anticoagulation | <input type="checkbox"/> Discharged | <input type="checkbox"/> Liver |
| <input type="checkbox"/> Aorta | <input type="checkbox"/> Diuretics | <input type="checkbox"/> Preload |
| <input type="checkbox"/> Clot | <input type="checkbox"/> Dressing | <input type="checkbox"/> Sepsis |
| <input type="checkbox"/> Complications | <input type="checkbox"/> Hygiene | <input type="checkbox"/> Sterile |
| <input type="checkbox"/> Consistent Diet | <input type="checkbox"/> Implant | <input type="checkbox"/> Sternotomy |
| <input type="checkbox"/> Dehydrated | <input type="checkbox"/> ICU | <input type="checkbox"/> Stroke |

Appendix

Scan the QR codes below to access additional reference materials.

My Important Numbers

Print this reminder checklist list off before each clinic visit checkup to keep all your current health measures and questions at your fingertips.

The form is titled "My Important Numbers" and includes sections for: "My HbA1c care team can be reached at:", "My HbA1c target value:" (with a field for 7.7), "My pump parameters: B15C, B15C, B15C, B15C, B15C, B15C", "I can feel my pump:" (Yes/No), "I take my blood glucose every 4 hours:" (Blood Pressure, Glucose, Weight), "My blood glucose goal:", "I am taking the following blood thinner(s):" (Coumadin, Other, Aspirin), "I have my strength changed:" (Yes/No), "My target weight:", "My daily step goal:", "My daily water intake goal:", and "Additional information:".



My Heart Success Medicines

Print this medicine tracker out and check off the medicines you are taking. Write (in pencil) any names or doses you need to remember since your heart medicines may change often. At your next visit, **ask your care team** if any of your medicine doses need to change.

The form is titled "My Heart Success Medicines" and includes a section for "I am on the following types of heart medicines:" with checkboxes for: ACE Inhibitor, ARB, Beta Blocker, Calcium Channel Blocker, Diuretic, Statin, and Other. It also has sections for "Diuretic", "Other Blood Pressure Medicine", and "Antiplatelet" with checkboxes for various medications like Hydrochlorothiazide, Lisinopril, Amlodipine, Aspirin, Clopidogrel, and Warfarin.



Appendix

My HeartMate 3™ LVAD Checklist

First seen in Chapter 3, download and keep this VAD maintenance milestone cheat sheet on hand to refer to often.

My HeartMate 3™ LVAD Checklist

Daily

- Recharge a compatible controller (i.e., Atrial battery) before each use to help and verify the light
- Before removing batteries from charger, allow batteries power before connecting to system controller
- Check that the air lines on the system controller are open
- Spin through 180 degrees and verify there is flow
- Check for the green light on the middle power unit before plugging in or replacing
- Check for leaks near the tubing back connecting to the main battery

Weekly

- Consider whether driving can be restricted
- Check the middle controller and check for any damage
- Check the battery charger in front of dust and debris
- Recharge the reserve batteries in the charger dock

Monthly

- Inspect and clean the battery and battery clip contacts with a cotton swab dampened with alcohol
- Check for any damage to connections
- Check the charger dock for any damage

3 Months - Yearly

- Replace the AC battery in the middle power unit
- Inspect both system controllers for damage
- Your caregiver will help you evaluate all of batteries. Check battery expiration date— if it has been used for more than 3 years, contact your care team
- If you are equipped with a pump care team, contact the back up controller to confirm repair and allow it to charge completely. Otherwise, perform a self-troubleshoot controller.

actionlivingnetwork.org | myheartmate.com **action** **Abbott**



VAD Emergency Contact Medical Safety Card

If you don't already have a medical safety card, download, print and fill out the ACTION card below (as seen in Chapter 7). Always carry your medical safety card with you in case of an emergency.

Medical Safety Card **action**
Living Network

Emergency Contact Information

If you don't already have a medical safety card, print this page and fill out the ACTION card below. See the sample card for more information on your emergency card. It is not valid for travel. It is an emergency card to be used in the event of an emergency. Always carry your medical safety card with you in case of an emergency.

First Name _____ <small>Print name in all caps and last name initials</small>	VAD Emergency Contact: _____ <small>Print name in all caps and last name initials</small>
Last Name _____ <small>Print name in all caps and last name initials</small>	XXX-XXX-XXXX _____ <small>Print phone number in all caps</small>
Address _____ <small>Print address in all caps</small>	Hospital Name _____ <small>Print hospital name in all caps</small>
City _____ <small>Print city in all caps</small>	XXXX Street Name, Apt _____ <small>Print street name in all caps</small>
State _____ <small>Print state in all caps</small>	City, State, Zip _____ <small>Print city, state, and zip in all caps</small>

First Name _____ **VAD Emergency Contact:** _____
Print name in all caps and last name initials

Last Name _____
Print name in all caps and last name initials

Address _____
Print address in all caps

City _____
Print city in all caps

State _____
Print state in all caps

City, State, Zip _____
Print city, state, and zip in all caps

actionlivingnetwork.org | myheartmate.com **action** **Living Network** **Abbott**



HeartMate Touch™ Communication System Overview: The HeartMate Touch™ Communication System is intended for use by clinicians in the hospital to wirelessly monitor a patient's HeartMate II™ Left Ventricular Assist System or HeartMate 3™ Left Ventricular Assist System. The HeartMate Touch Communication System is required during implant procedures and any time close monitoring of system operation is needed. It provides clinicians with the ability to program system parameters such as pump speed, assess and track alarm conditions, and view and save performance data.

Abbott

6101 Stoneridge Dr., Pleasanton, CA 94588 USA, Tel: 1 925 847 8600
Cardiovascular.Abbott/HeartMate3

Rx Only

Brief Summary: Prior to using these devices, please review the Instructions for Use for a complete listing of indications, contraindications, warnings, precautions, potential adverse events and directions for use.

HeartMate 3™ LVAS Indications: The HeartMate 3™ Left Ventricular Assist System is indicated for providing short- and long-term mechanical circulatory support (e.g., as bridge to transplant or myocardial recovery, or destination therapy) in adult and pediatric patients with advanced refractory left ventricular heart failure and with an appropriate body surface area.

HeartMate II™ LVAS Indications: The HeartMate II™ Left Ventricular Assist System is indicated for use as a "bridge to transplantation" for cardiac transplant candidates who are at risk of imminent death from non-reversible left ventricle failure. It is also indicated for use in patients with New York Heart Association (NYHA) Class IIIB or IV end-stage left ventricular failure, who have received optimal medical therapy for at least 45 of the last 60 days, and who are not candidates for cardiac transplantation. The HeartMate II Left Ventricular Assist System is intended for use both inside and outside of the hospital, or for transportation of Left Ventricular Assist Device patients via ground ambulance, airplane, or helicopter.

HeartMate 3™ and HeartMate II™ LVAS Contraindications: The HeartMate 3 and HeartMate II Left Ventricular Assist Systems are contraindicated for patients who cannot tolerate, or who are allergic to, anticoagulation therapy.

HeartMate 3™ and HeartMate II™ LVAS Adverse Events: Adverse events that may be associated with the use of the HeartMate 3 or HeartMate II Left Ventricular Assist System are listed below: death, bleeding, cardiac arrhythmia, localized infection, right heart failure, respiratory failure, device malfunctions, driveline infection, renal dysfunction, sepsis, stroke, other neurological event (not stroke-related), hepatic dysfunction, psychiatric episode, venous thromboembolism, hypertension, arterial non-central nervous system (CNS) thromboembolism, pericardial fluid collection, pump pocket or pseudo pocket infection, myocardial infarction, wound dehiscence, hemolysis (not associated with suspected device thrombosis) and possible pump thrombosis.

™ Indicates a trademark of the Abbott group of companies.

© 2024. All Rights Reserved.

MAT-2405453 v1.0 | Item approved for U.S. use only.

actionlearningnetwork.org | myactioneducation.org

 @actionlearningnetwork  @Action4HF

06.2024

action 

 **Abbott**